

**Tuesday, April 7, 2020 at 6:30 PM ET**

**The Capital Grille**

5197 Big Island Drive, Jacksonville, FL 32246

**Speaker: David Kostick, MD**

**RSVP Jason Ambrose**

**JAmbrose@horizontherapeutics.com**

**By April 3, 2020**

**(904) 704-4466**

### Program Highlights

The objectives of this program are to:

- Provide an overview of Thyroid Eye Disease (TED) and review the pathophysiology
- Describe the mechanism of action of TEPEZZA, a new treatment option for TED
- Present clinical data highlighting the efficacy and safety of TEPEZZA for TED
- Describe how to administer, monitor, and manage potential adverse reactions for TEPEZZA

### INDICATION

TEPEZZA is indicated for the treatment of Thyroid Eye Disease.

### IMPORTANT SAFETY INFORMATION

#### Warnings and Precautions

**Infusion Reactions:** TEPEZZA may cause infusion reactions. Infusion reactions have been reported in approximately 4% of patients treated with TEPEZZA. Reported infusion reactions have usually been mild or moderate in severity. Signs and symptoms may include transient increases in blood pressure, feeling hot, tachycardia, dyspnea, headache, and muscular pain. Infusion reactions may occur during an infusion or within 1.5 hours after an infusion. In patients who experience an infusion reaction, consideration should be given to premedicating with an antihistamine, antipyretic, or corticosteroid and/or administering all subsequent infusions at a slower infusion rate.

**Preexisting Inflammatory Bowel Disease:** TEPEZZA may cause an exacerbation of preexisting inflammatory bowel disease (IBD). Monitor patients with IBD for flare of disease. If IBD exacerbation is suspected, consider discontinuation of TEPEZZA.

**Hyperglycemia:** Increased blood glucose or hyperglycemia may occur in patients treated with TEPEZZA. In clinical trials, 10% of patients (two-thirds of whom had preexisting diabetes or impaired glucose tolerance) experienced hyperglycemia. Hyperglycemic events should be managed with medications for glycemic control, if necessary. Monitor patients for elevated blood glucose and symptoms of hyperglycemia while on treatment with TEPEZZA. Patients with preexisting diabetes should be under appropriate glycemic control before receiving TEPEZZA.

#### Adverse Reactions

The most common adverse reactions (incidence  $\geq 5\%$  and greater than placebo) are muscle spasm, nausea, alopecia, diarrhea, fatigue, hyperglycemia, hearing impairment, dysgeusia, headache, and dry skin.

For additional information on TEPEZZA, please see [Full Prescribing Information](#) at [TEPEZZAhcp.com](#).

Please note that Horizon Therapeutics adheres to the principles set forth in the PhRMA Code on Interactions with Health Care Professionals. All program functions are for invited guests only. Spouses and others, including office staff or other family members, may not attend.

REPORTING DISCLOSURE: All meal participants will be required to sign-in at this event. As required by the federal Sunshine Act and other similar state laws, the cost of the meal provided to you will be publicly reported. If you want to attend the program without your information reported, you may opt-out of the meal.